## Medicine and Doses for Pain and Fever for Children Under Age 2





Follow these steps to be sure you give the right dose (amount of medicine) to your baby.

- **Step 1:** Look at the bottle and find the name of the medicine. Use Table 1 for acetaminophen and Table 2 for ibuprofen.
- **Step 2:** Next, find the concentration of the medicine on the label. Match to the concentration listed in the table.
- **Step 3:** Move down the column on the left of the table to find your child's weight. (*Do not give ibuprofen to a baby less than 6 months old.*)
- **Step 4:** Find the column on the right that matches the concentration of your medicine.
- **Step 5:** Always use a syringe or dosing cup to measure your child's medicine.

## **Table 1. Acetaminophen (Tylenol or another brand)**

\* please note the concentration of medicine on the label of your bottle






Give every 4 to 6 hours, as needed, no more than 5 times in 24 hours			
Child's Weight	80 mg in each 0.8 mL (Infants' Concentrated Drops)	<b>160 mg in each 5 mL</b> (Infants' or Children's Liquid)	
6 to 11 pounds (2.7 to 5 kilograms)	0.4 mL	1.25 mL	
12 to 17 pounds (5.5 to 7.7 kilograms)	0.8 mL	2.5 mL	
18 to 23 pounds (8.2 to 10.5 kilograms)	1.2 mL	3.75 mL	
24 pounds (10.9 kilograms) or more	Follow directions on medicine package		

## Table 2. Ibuprofen (Motrin, Advil, or another brand)

\* please note the concentration of medicine on the label of your bottle



Infant or children's ibuprofren liquid:		
100mg/5mL		

Give every 6 to 8 hours, as needed			
Child's Weight	50 mg in each 1.25 mL (Infants' Concentrated Drops)	<b>100 mg in each 5 mL</b> (Infants' or Children's Liquid)	
less than 11 pounds (5 kilograms) or less than 6 months old	DO NOT GIVE	DO NOT GIVE	
12 to 17 pounds (5.5 to 7.7 kilograms)	1.25 mL	2.5 mL	
18 to 23 pounds (8.2 to 10.5 kilograms)	1.875 mL	3.75 mL	
24 pounds (10.9 kilograms) or more	Follow directions on medicine package		